



Food Additives & Contaminants: Part B Surveillance

ISSN: 1939-3210 (Print) 1939-3229 (Online) Journal homepage: <http://www.tandfonline.com/loi/tfab20>

Pesticide residues in leafy vegetables and human health risk assessment in North Central agricultural areas of Chile

Sebastian Elgueta, Stella Moyano, Paulina Sepúlveda, Carlos Quiroz & Arturo Correa

To cite this article: Sebastian Elgueta, Stella Moyano, Paulina Sepúlveda, Carlos Quiroz & Arturo Correa (2017): Pesticide residues in leafy vegetables and human health risk assessment in North Central agricultural areas of Chile, Food Additives & Contaminants: Part B, DOI: [10.1080/19393210.2017.1280540](https://doi.org/10.1080/19393210.2017.1280540)

To link to this article: <http://dx.doi.org/10.1080/19393210.2017.1280540>



Accepted author version posted online: 16 Jan 2017.
Published online: 29 Jan 2017.



Submit your article to this journal [↗](#)



Article views: 8



View related articles [↗](#)



View Crossmark data [↗](#)

ARTICLE

Pesticide residues in leafy vegetables and human health risk assessment in North Central agricultural areas of Chile

Sebastian Elgueta^a, Stella Moyano^a, Paulina Sepúlveda^b, Carlos Quiroz^c and Arturo Correa^a

^aDepartment of Environment and Sustainability, Instituto de Investigaciones Agropecuarias, Santiago, Chile; ^bDepartment of Plant Pathology, Instituto de Investigaciones Agropecuarias, Santiago, Chile; ^cDepartment of Plant Health, Instituto de Investigaciones Agropecuarias, La Serena, Chile

ABSTRACT

To investigate pesticide residue concentrations and potential human health risk, a study was conducted in 118 leafy vegetable samples collected in 2014–2015 from the North Central agricultural areas of Chile. The pesticide residues were determined using the multiresidue QuEChERS method by gas chromatography as well as high-performance liquid chromatography. The results indicated that 27% of the total samples contained pesticide residues above the maximum residue limits of each active ingredient. The maximum estimated daily intake obtained for carbon disulphide (CS₂), methamidophos, azoxystrobin and cypermethrin were 0.57, 0.07, 0.06 and 0.05 mg kg⁻¹, respectively, which was higher than their acceptable daily intake. It is concluded that inhabitants of the North Central agricultural area of Chile are not exposed to health risks through the consumption of leafy vegetables with the exception of methamidophos. Nevertheless, the high levels of methamidophos detected in leafy vegetables could be considered a potential chronic health risk.

ARTICLE HISTORY

Received 5 October 2016
Accepted 3 January 2017

KEYWORDS

Pesticide residues; maximum residue limits (MRLs); acceptable daily intake (ADI); hazard quotient (HQ)

Introduction

The demand for food is increasing worldwide because of the high global population and improvement of living standards. The use of pesticides in modern agriculture is necessary for most crops to guarantee the food supply. However, improper and excessive use of pesticides in horticultural processes has resulted in environmental pollution and increasing human health risk (Oerke & Dehne 2004; Hernández et al. 2013; Köck-Schulmeyer et al. 2013). The World Health Organization (WHO) promotes the consumption of vegetables and fruits as an important source of essential vitamins, minerals and antioxidants. This food group has active compounds that can reduce the effect of cardiovascular diseases (Ivey et al. 2015). However, this food group can be a source of contamination, because the ingestion of pesticide residues is a potential source of diseases and negative effects (Parrón et al. 2014; Lemos et al. 2016).

It is reasonable to assume that vegetables and fruits may contain pesticide residues in harmless concentrations (Chen et al. 2012; Li et al. 2015). The maximum residue limits (MRLs), expressed as the highest concentration of pesticide residues in vegetables and fruits (mg kg⁻¹), are used to determine the maximum intake legally permitted by regulations introduced by different

countries and internationally by the Codex Alimentarius Commission (Darko & Akoto 2008; CODEX Alimentarius Commission 2011; Gad Alla et al. 2015). The human intake of pesticide residues in vegetables can be much higher than the intake related to water consumption and air inhalation. It is very important to monitor such chemicals in vegetables and to assess their potential risks to human health (Szyrka et al. 2015). Some indicators can estimate the amount of chemicals in food, which is represented as the acceptable daily intake (ADI) and expressed as mg kg⁻¹ body weight (bw) day⁻¹. The ADI can be used to predict the daily intake during a lifetime that would prevent health risks to the consumer (FAO/WHO 1997; Akoto et al. 2013) and is used to determine the estimated daily intakes (EDIs).

The aim of obtaining data on pesticide residue in food is to improve pest management programmes (Bhanti & Taneja 2005). Most organophosphate pesticides such as methamidophos, parathion-methyl, parathion, monocrotophos and phosphamidon are toxic. Therefore, numerous types have been banned since 2007 for agricultural use in China (Chen et al. 2011; Yu et al. 2016). In Brazil, the registration of methamidophos was recently cancelled and the phaseout period ended in July 2012 (Jardim et al. 2014). In some countries

including Guatemala, Spain, Jamaica and China, the use of methamidophos is still allowed but adverse effects have been detected (Glat 2010). In Chile, there are no international scientific articles reporting the monitoring of pesticide residue in food and their health risk assessment. The lack of information on the health risk associated with the exposure of the population to pesticides through consumption of vegetables is of important concern. Since 2009, Chile has been part of the Organisation for Economic Co-operation and Development (OECD) that promotes policies to improve the economic and social well-being of people. The statistical analysis by the OECD reports that pesticides are used at a total average of 0.21 kg ha⁻¹. Chile has one of the highest levels of pesticide use with a total of 10.7 kg ha⁻¹ used in 2009. The high level of pesticide use can be explained by the severe phytosanitary food certification standards that international markets have imposed on Chile (Engler et al. 2012; Handschuch et al. 2013; Melo et al.).

Vegetables are produced in all regions in Chile, with nearly 34,000 farms, 65% of which have less than 5 ha, indicating that this sector is composed mainly of small growers (INE 2002, 2016). The National Institute of Statistics in Chile estimated that the national horticultural area consisted of almost 70,000 ha by 2014, 84% of which is between the regions of Coquimbo and Maule (30°00' S–72°30' W to 36°05' S–72°30' W). Leafy vegetables such as lettuce (*Lactuca sativa* L.), chard (*Beta vulgaris* subsp. *vulgaris*) and spinach (*Spinacia oleracea*) are among the main leafy vegetables produced in Chile. The production of lettuce in Chile has been estimated at over 75,000 tons and is concentrated (74%) mainly in the Valparaíso (33°2' S–71°40' W) to the Metropolitana (33°27' S–70°40' W) regions (Oficina de Estudios y Políticas Agrarias). The purposes of this study were (1) to quantify the concentration of pesticide residues in leafy vegetables in three regions of Chile and (2) to assess the health risks associated with the consumption of potentially contaminated vegetables. The relevance of this work is to enhance the understanding of the importance of pesticide residues in food and its effects on human health, which would improve the methods of pesticide use in integrated pest management programmes in Chile.

Materials and methods

Study area

The Coquimbo, Valparaíso and Metropolitana regions are located in North Central Chile, which is the most important growing area for leafy vegetables. The local climate is characterised by short, rainy winters and warm summers. The average annual temperature

ranges from 12°C to 14°C, whereas the average annual rainfall is from 78 mm in Coquimbo to 312 mm in the Metropolitan Region. More than 75% of the rainfall occurs in June, July and August.

Leafy vegetable sampling

To conduct a representative study, it was necessary to determine the appropriate sample size (n) based on the number of leafy vegetable (lettuce, chard and spinach) growers in the area under study (Table 1), using the following formula (INE 2002, 2016):

$$n = (N \times Z^2 \times p \times q) / (d^2 \times [N-1] + Z^2 \times p \times q)$$

where N is the total number of growers, Z^2 is 1,962 (security 95%), p is the expected proportion (5% = 0.05), $q = 1 - p$ (1 - 0.05 = 0.95) and d is the precision (3%). A total of 118 samples of leafy vegetables consisting of lettuce (74), chard (26) and spinach (18) from all regions were taken and analysed during the 2014–2015 cultivation season. The sample size was 2 kg of leaves collected from the sites, wrapped in aluminium foil and transported to the laboratory. Samples were milled and homogenised, 20 g were transferred into 100 mL glass containers and stored at –20°C until the analysis.

Chemicals and reagents

Analytical grade pesticide standards (99% purity) were obtained from Chem Service (West Chester, PA, USA) and Pestanal (Sigma-Aldrich, Santiago, Chile). The standard working solutions were prepared by diluting the stock solutions with acetone/isooctane. All analytical reagents were purchased from Merck and J.T. Baker (Santiago, Chile). All the solutions were prepared with ultra-high purity water and filtered using a Milli-Q system from Millipore (IADET, Santiago, Chile).

Multiresidue method and instrumentation

The pesticides analysed in this study are the most frequently detected and used in pest management

Table 1. Number of growers of leafy vegetables in North Central areas of Chile (INE 2002, 2016).

| Regions | Leafy vegetables | | |
|---------------|------------------|---------|---------|
| | Chard | Spinach | Lettuce |
| Coquimbo | 46 | 7 | 263 |
| Valparaíso | 160 | 128 | 767 |
| Metropolitana | 210 | 60 | 483 |
| Total | 416 | 195 | 1513 |

programmes by growers in Chile. The MRLs in Chile were established by the RES-33/2010-02-16 and RES-76272011-10-02 based on the MRLs of Codex Alimentarius, the European Union (EU) and the USA. All pesticide residues present in the leafy vegetables were determined using gas chromatography-mass spectrometry (GC-MS). The standard method EN 15,662 of the European Committee for Standardization for food material of plant origin, known as the quick, easy, cheap, effective, rugged and safe (QuEChERS) method, was used in this study. This method was previously described by Anastassiades et al. (2003) and Payá et al. (2007). Two tubes in the QuEChERS system Waters (Milford, CT, USA) were used for extraction and clean-up. The first tube contained 4 g magnesium sulphate ($MgSO_4$), 1 g sodium chloride (NaCl) and 1.5 g citrate. The second tube contained 900 mg $MgSO_4$, 150 mg primary secondary amine and 150 mg C18. Briefly, 10 g of each homogenised leafy vegetable sample was weighed, placed into a 50-mL centrifuge tube and extracted with 10 mL acetonitrile with vigorous shaking for 1 min. After 3 min, the mixture was centrifuged at 2500 rpm for 5 min and the total acetonitrile supernatant phase was transferred to a new 15-mL tube for the clean-up. The tube was shaken vigorously for 1 min and then centrifuged for 1 min at 2500 rpm. Then 6 mL of the supernatant was placed in a glass vial with ethyl acetate acidified with 100 μ L formic acid, evaporated to dryness, re-dissolved in 1 mL acetone/isooctane (1:9) and filtered through a 0.20 μ m polyvinylidene fluoride (PVDF) syringe filter into an autosampler vial for the analysis.

Organophosphate residues were quantified using an Agilent 7890-Autosampler (Wilmington, DE, USA) gas chromatography system with a nitrogen-phosphorus detector (NPD) detector. The GC system was equipped with a fused silica capillary column HP-5MS (30 m \times 0.25 mm \times 0.25 μ m; Agilent, Santa Clara, CA, USA). For the halogenated pesticides, a GC-electron capture detector (ECD) and a Perkin Elmer auto-system XL (Waltham, MA, USA) were used. The GC-ECD system was equipped with a fused silica capillary column (DB-1 and DB-17, 30 m \times 0.25 mm \times 0.25 μ m; Sigma-Aldrich (St. Louis, MO, USA)). The methyl-carbamates were quantified using a high-performance liquid chromatography (HPLC) system with a Merck Hitachi LaChrom D-7000-autosampler (Dartford, United Kingdom) with ultraviolet (UV) and fluorescence (FL) detectors and a 655A-B reaction pump. All samples were filtered through a 0.22- μ m PVDF syringe filter into an autosampler vial before analysis. The HPLC was equipped with a UV detector set to 270 nm and a column (RP-18 X-Terra, 3.9 \times 150 mm; Waters). The mobile phase was

acetonitrile/water (30/70, % v/v). The samples were filtered through a 0.22- μ m PVDF syringe filter into an autosampler vial before analysis. For imidacloprid and carbendazim, an HPLC system with a Merck Hitachi D-6000 (Burladingen, Germany) was used. The dithiocarbamate levels were determined using distillation and quantification with a Thermo-Genesys 10VIS detector (Madison, WI, USA), using the method of Perz et al. (2000). The results were expressed as milligrams of carbon disulphide per kilogram ($mg CS_2 kg^{-1}$).

Method performance and quantification

The methods were validated according to the SANCO document 12571/2013 guidelines. The limit of quantification (LOQ), which was defined as the lowest concentration that could be quantified with an acceptable recovery (70–105%), started at 0.01 $mg kg^{-1}$ and was further increased depending on the pesticide (Table 2). The precision, which was expressed as the relative standard deviation (RSD, %) was determined by analysing replicate samples at three levels ($n = 5/\text{level}$) and was set at <20%. The average recovery of the different pesticides varied between 70% and 105%. The limit of detection (LOD) was calculated according to the Environmental Protection Agency recommendations. The quantification was performed using a mixture containing known amounts of the pesticides and the detector response for each compound was determined. The area of the corresponding peak in the sample was compared with that of the known standard (Bempah et al. 2012; Gad Alla et al. 2015).

Human health risk assessment

Pesticide residue concentrations on the leafy vegetables were determined as the arithmetic mean of all results obtained. The EDI was calculated by multiplying the pesticide concentration of each pesticide ($mg kg^{-1}$) with the food consumption rate ($kg day^{-1}$) and dividing this by the bw (Gad Alla et al. 2015). Food consumption was based on data from the WHO/Global Environment Monitoring (GEMS)/Food Cluster Diet 2012, a part of the FOSCOLLAB, the WHO Global platform for food safety data and information. Cluster G05 for Chile reports a leafy vegetable consumption of 2.046 $g day^{-1}$. The average bw used for Chileans was defined as 72.3 kg (ENS Chile 2009-2010). The EDI ($mg kg^{-1} bw day^{-1}$) was calculated as follows: $EDI = \sum RL_i \times F_i/bw$; where RL_i is the residue level of leafy vegetable i and F_i is the food consumption rate of food item i .

The hazard quotient (HQ) was considered as the potential risk of adverse health effects from a mixture of pesticides to indicate the long-term risk assessment and

Table 2. Validation parameters ($n = 5$) for QuEChERS multiresidue method at three levels in leafy vegetables and MRLs.

| Pesticide | Recovery % | RSD % | LOQ (mg kg ⁻¹) | LOD (mg kg ⁻¹) | MRL (mg kg ⁻¹) | Instrument | |
|----------------------------|------------|-------|----------------------------|----------------------------|----------------------------|-------------------------------|---------|
| Chlorothalonil | 75.7 | 13.4 | 0.01 | 0.005 | 0.01 | GC-ECD | |
| Chlorpyrifos | 92.5 | 1.1 | 0.01 | 0.005 | 0.05 | | |
| Difenoconazole | 103.6 | 7.1 | 0.01 | 0.005 | 0.05 | | |
| Lambda-cyhalothrin | 98.0 | 3.5 | 0.01 | 0.005 | 0.05 | | |
| Biphenethrin | 92.8 | 3.2 | 0.01 | 0.005 | 0.05 | | |
| Boscalid | 99.8 | 3.6 | 0.01 | 0.005 | 5 | | |
| Cyfluthrin | 92.6 | 3.1 | 0.02 | 0.01 | 0.02 | | |
| Deltamethrin | 95.0 | 7.8 | 0.01 | 0.005 | 0.02 | | |
| Cypermethrin | 93.6 | 8.1 | 0.01 | 0.005 | 0.7 | | |
| Azoxystrobin | 94.9 | 3.0 | 0.01 | 0.005 | 0.01 | | |
| Esfenvalerate | 89.9 | 3.7 | 0.02 | 0.005 | 0.02 | | |
| Endosulfan | 91.4 | 3.3 | 0.01 | 0.005 | 0.05 | | |
| Linuron | 93.4 | 12.7 | 0.01 | 0.005 | 0.01 | | |
| Permethrin | 91.4 | 0.8 | 0.01 | 0.005 | 2 | | |
| Propiconazole | 103.4 | 4.6 | 0.01 | 0.005 | 0.05 | | |
| Azinphos-methyl | 100.0 | 4.0 | 0.01 | 0.005 | 0.5 | | |
| Metalaxyl | 96.1 | 2.2 | 0.01 | 0.005 | 0.05 | | |
| Methamidophos | 89.3 | 4.4 | 0.01 | 0.005 | 0.01 | | |
| Diazinon | 86.3 | 5.4 | 0.01 | 0.005 | 0.5 | | |
| Dimethoate | 103.0 | 5.9 | 0.01 | 0.005 | 0.02 | | |
| Fludioxonil | 100.7 | 4.5 | 0.01 | 0.005 | 0.05 | | |
| Phosmet | 103.2 | 4.2 | 0.01 | 0.005 | 0.05 | | |
| Malathion | 95.7 | 3.2 | 0.01 | 0.005 | 0.02 | GC-NPD | |
| Methidathion | 93.9 | 2.9 | 0.01 | 0.005 | 0.02 | | |
| Penconazole | 90.4 | 4.1 | 0.01 | 0.005 | 0.05 | | |
| Pyrimethanil | 101.7 | 3.1 | 0.01 | 0.005 | 0.05 | | |
| Pyraclostrobin | 87.6 | 7.1 | 0.01 | 0.005 | 0.5 | | |
| Thiabendazole | 95.3 | 5.6 | 0.01 | 0.005 | 0.05 | | |
| Thiamethoxam | 92.2 | 3.0 | 0.01 | 0.005 | 0.01 | | |
| Triadimefon | 97.4 | 0.5 | 0.01 | 0.005 | 0.1 | | |
| Abamectin | 104.3 | 5.8 | 0.002 | 0.001 | 0.01 | | HPLC-FL |
| Aldicarb | 86.9 | 4.4 | 0.01 | 0.005 | 0.02 | | |
| Methomyl | 94.4 | 5.1 | 0.01 | 0.005 | 0.01 | | |
| CS ₂ (mancozeb) | 91.8 | 5.3 | 0.01 | 0.005 | 0.05 | Spectro-photometer HPLC-UV | |
| Carbendazim | 70.0 | 1.57 | 0.02 | 0.01 | 0.05 | | |
| Imidacloprid | 95.3 | 1.3 | 0.02 | 0.01 | 0.05 | | |

was calculated by dividing the EDI by the relevant ADI. The ADI values are given by the European Commission and the European Food Safety Authority. HQ was calculated using the formula $HQ = EDI/ADI \times 100$, a percentage >100 indicating that the exposed people were unlikely to experience obvious toxic effects. Furthermore, an index below 100 indicated the possibility that the exposure could induce toxic effects (Yu et al. 2016). HQ was calculated for the pesticides and leafy vegetables and the results were summed up to obtain a chronic hazard index (cHI) where, $cHI = \sum HQ$. $cHI > 100$ indicated that the leafy vegetables should be considered a risk to the consumers, whereas an index below 100 indicated that the consumption of the leafy vegetable was considered acceptable (Gad Alla et al. 2015).

Results and discussion

Analytical results

The accuracy of the QuEChERS method was expressed as recovery (%) and RSD (%) at the levels 0.01, 0.1 and

0.5 mg kg⁻¹ in leafy vegetables. The recovery values varied at 70–105%. These results are concordant with 70–120% as given in the EU guidance document on pesticides analysis (SANTE, European Commission 2016). In addition, the RSD obtained was 0.5–13.4%, which is lower than the 15% required to validate extraction and clean-up (WHO/FAO 2013). The LOD and LOQ for the analytical method varied at 0.005–0.02 mg kg⁻¹ for the leafy vegetables. Similar results were previously obtained by Srivastava et al. (2011) for 20 vegetables including leafy samples. They extracted their samples using the QuEChERS method, measured with a GC-ECD/NPD and obtained a recovery of 70–96% and RSD of 15% for the 35 pesticides analysed.

Table 3 gives an overview of the obtained occurrence data. Pesticide residues were found in 27, 9 and 5 samples of lettuce (36%), chard (35%) and spinach (72%), respectively. A mixture of pesticide residues was shown in 47, 17 and 13 samples of lettuce (64%), chard (65%) and spinach (72%). Among the leafy vegetables, 27% of the total samples contained pesticide residues above the MRLs of each active ingredient, whereas 42%,

Table 3. Pesticide residues in leafy vegetables in North Central agricultural areas in Chile. All samples above LOD contained multiple residues.

| Vegetable | <i>n</i> | <i>n</i> < LOD | | <i>n</i> > LOD | | <i>n</i> > MRL | |
|-----------|----------|----------------|-----|----------------|-----|----------------|-----|
| Lettuce | 74 | 27 | 36% | 47 | 64% | 16 | 22% |
| Chard | 26 | 9 | 35% | 17 | 65% | 11 | 42% |
| Spinach | 18 | 5 | 28% | 13 | 72% | 5 | 28% |
| total | 118 | 41 | 35% | 77 | 65% | 32 | 27% |

28% and 22% of the pesticides detected were above the MRLs for chard, spinach and lettuce, respectively. In brief, two or more pesticide residues were detected in 65% of the samples. These results can be attributed to the high frequency of pesticide application on farms, the short time intervals allowed between the numerous applications (McEwen et al. 1980) and the pesticide retention capacity of the leafy vegetable surfaces (Li et al. 2014; Gad Alla et al. 2015; Yu et al. 2016).

These results are in accordance with those of Yu et al. (2016) who monitored organophosphorus pesticides (OPs) in leafy vegetables in Changchun, China. In that study, only 7% of the total samples were free from pesticide residues, whereas 23% and 68% of the total samples contained OPs above and below the MRLs, respectively. In addition, González-Rodríguez et al. (2008) monitored pesticide residues in 75 leafy vegetable samples (chard, spinach and lettuce). They reported that pesticide residues were above the MRLs in 15 of 75 analysed samples. The highest concentrations of the insecticides procymidone and cypermethrin (12 and 6 mg kg⁻¹) were found in lettuce and chard, respectively. Esturk et al. (2011) detected a high occurrence of

pesticide residues in 120 lettuce and spinach samples, most of which were not authorised for use in Turkey.

Occurrence and levels

Table 4 illustrates the occurrence and concentrations of pesticide residues in the analysed samples. Methamidophos, chlorpyrifos, boscalid and lambda-cyhalothrin were the most predominant compounds found in all the samples, followed by carbendazim and imidacloprid, which were found in lettuce and chard. Methamidophos showed the highest concentration (29.47 mg kg⁻¹), followed by chlorpyrifos (6.86 mg kg⁻¹), which were detected in lettuce, chard and spinach. The levels of methamidophos found in this study could induce serious adverse effects on human health and the environment. Goh et al. (1990) provided the first evidence that people could be affected by consumption of leafy vegetables with high levels of methamidophos and profenofos, as they showed that blood cholinesterase levels of hospitalised patients were depressed by 26–81%.

The results showed that cypermethrin, cyfluthrin, difenoconazole, esfenvalerate, metalaxyl, permethrin and thiamethoxam were found in 1% of the samples at values below the MRLs. However, metalaxyl exceeded the MRL for chard (0.13 mg kg⁻¹). Moreover, boscalid, chlorothalonil, difenoconazole and metalaxyl found in the chard samples are not authorised for use in this vegetable. Their presence could be attributed to a wrong pesticide choice made by the farmers for pest

Table 4. Pesticide residues in leafy vegetables in North Central agricultural areas in Chile.

| Active substance | Category | <i>N</i> > LOD/frequency | | Min-max (mg kg ⁻¹) | Mean (mg kg ⁻¹) | MRLs Chile (mg kg ⁻¹) | |
|--------------------|----------|--------------------------|-----|--------------------------------|-----------------------------|-----------------------------------|---------------|
| Cypermethrin | I | 1 | 1% | 0–1.61 | 1.61 | 0.70 | (L, C, S) |
| Azoxystrobin | F | 3 | 3% | 0.39–3.38 | 2.25 | not available | not available |
| Boscalid | F | 15 | 13% | 0.01–3.06 | 0.52 | 30.00 | (L, C, S) |
| Carbendazim | F | 13 | 11% | 0.01–1.38 | 0.20 | 0.10 | (L, C, S) |
| Cyfluthrin | I | 1 | 1% | 0–0.48 | 0.48 | 3.00 | (L, C, S) |
| Chlorothalonil | F | 2 | 2% | 0.11–1.42 | 0.77 | 0.01 | (L, C, S) |
| Chlorpyrifos | I | 16 | 14% | 0.01–6.86 | 0.54 | 0.05 | (L, C, S) |
| CS2 (Mancozeb) | F | 5 | 4% | 0.34–4.0 | 20.32 | 10.00 | (L) |
| | F | 4 | 3% | 0.38–5.88 | 2.20 | 0.05 | (S) |
| Difenoconazole | F | 1 | 1% | 0–0.08 | 0.08 | 0.05 | (C) |
| | F | 5 | 4% | 0.32–3.14 | 0.95 | 2.00 | (L) |
| Esfenvalerate | I | 1 | 1% | 0–0.3 | 0.30 | 0.02 | (L, C, S) |
| Imidacloprid | I | 6 | 5% | 0.02–0.18 | 0.11 | 3.50 | (L) |
| | I | 4 | 3% | 0.02–1.07 | 0.52 | 0.05 | (C) |
| Lambda-cyhalothrin | I | 14 | 12% | 0.029–1 | 0.22 | 2.00 | (L) |
| | I | 5 | 4% | 0.01–0.80 | 0.53 | 0.50 | (C) |
| Metalaxyl | F | 1 | 1% | 0–0.13 | 0.13 | 0.05 | (C) |
| | F | 5 | 4% | 0.02–1.76 | 0.50 | 2.00 | (L) |
| Methamidophos | I | 30 | 25% | 0.01–29.47 | 2.61 | 0.01 | (L, C, S) |
| Methomyl | I | 3 | 3% | 0.01–0.02 | 0.03 | 0.20 | (L, C, S) |
| Permethrin | I | 1 | 1% | 0–1.45 | 1.45 | 20 | (L, C, S) |
| Pyraclostrobin | F | 2 | 2% | 0.31–1.25 | 0.78 | 29.00 | (L, C, S) |
| Propiconazole | F | 2 | 2% | 0.03–0.05 | 0.04 | 0.05 | (L, C, S) |
| Thiamethoxam | I | 1 | 1% | 0–0.71 | 0.71 | 5 | (L) |

I: insecticide; F: fungicide; L = lettuce, C: chard, S: spinach.

control in chard or wrong advice from pesticide retailers (Wang et al. 2013). Furthermore, boscalid was also detected in spinach and lettuce, although it is an unauthorised active ingredient for these vegetables. Similar results have also been found in Thailand and China (Sapbamrer & Hongsihsong 2014; Yu et al. 2016).

The results also showed the presence of methamidophos residues in lettuce, chard and spinach with a frequency of 25% and a mean of 2.61 mg kg⁻¹, which exceeded the MRLs for leafy vegetables. Similarly chlorpyrifos, detected at a frequency of 14% and mean of 0.54 mg kg⁻¹ and carbendazim, showing a frequency of 11% and a mean of 0.20 mg kg⁻¹, in both cases exceeding the MRLs. Mancozeb, difenoconazole, lambda-cyhalothrin and thiamethoxam exceeded the MRLs for lettuce, whereas only difenoconazole and metalaxyl exceeded the MRLs in chard. The sum of the mean pesticide residues was highest in lettuce, followed by chard and spinach.

Similar situations to those observed in our study have been described due to the use of unauthorised or a high amount of pesticides, leading to high residue levels. De Putter and Van Sauers-Muller (2008) presented several reports of values as high as 1800 times the MRLs for methamidophos in spinach over the period 2004–2005. In addition, Akoto et al. (2015) found that methamidophos exceeded the MRLs in other commodities.

Human risk assessment

As shown in Table 5, the maximum EDI was obtained for mancozeb, methamidophos, azoxystrobin and cypermethrin at 0.57, 0.07, 0.06 and 0.05 mg kg⁻¹, which is 11-, 73-, 30- and 0.3-fold, respectively, higher than the corresponding ADI values. Freeman et al. (2016) found methamidophos exceedances of the ADI at the 25th percentile in vegetables in Israel, demonstrating potential exposure of the population to pesticide residues. Park et al. (2016) revealed for leafy vegetables in South Korea EDI and ADI exceedances of the MRLs in a range of 0.003–30%. Similar results were obtained by Chen et al. (2011), who detected cypermethrin in 18.7% of the investigated vegetable samples and EDI values ranging from 0.1% of the ADI for cyfluthrin to 2.61% of the ADI for omethoate. Yu et al. (2016) found methamidophos and phorate to be key components contributing to the potential health risks of inhabitants of the agricultural areas in China.

HQs for methamidophos, cypermethrin, mancozeb, cyfluthrin and lambda-cyhalothrin were 73.9, 30.4, 11.5, 4.5 and 3.0, respectively. HQs decreased in the order: methamidophos > cypermethrin > cyfluthrin > chlorpyrifos > chlorothalonil > pyraclostrobin > esfenvalerate >

Table 5. Estimated daily intake (mg kg⁻¹ bw day⁻¹) and potential health risk of pesticide residues in leafy vegetables.

| Pesticide | ADI | EDI | HQ |
|------------------------|-------|------|------|
| Cypermethrin | 0.002 | 0.05 | 30.4 |
| Azoxystrobin | 0.200 | 0.06 | 0.3 |
| Boscalid | 0.040 | 0.01 | 0.4 |
| Carbendazim | 0.080 | 0.01 | 0.1 |
| Cyfluthrin | 0.003 | 0.01 | 4.5 |
| Chlorothalonil | 0.015 | 0.02 | 1.4 |
| Chlorpyrifos | 0.010 | 0.02 | 1.5 |
| CS2 (mancozeb) L | 0.050 | 0.57 | 11.5 |
| CS2 (mancozeb) S | 0.050 | 0.06 | 1.2 |
| Difenoconazole (C) | 0.010 | 0.00 | 0.2 |
| Difenoconazole (L) | 0.010 | 0.03 | 2.7 |
| Esfenvalerate | 0.020 | 0.01 | 0.4 |
| Imidacloprid (L) | 0.060 | 0.00 | 0.0 |
| Imidacloprid (C) | 0.060 | 0.01 | 0.2 |
| Lambda-cyhalothrin (L) | 0.005 | 0.01 | 1.2 |
| Lambda-cyhalothrin (C) | 0.005 | 0.02 | 3.0 |
| Metalaxyl (C) | 0.080 | 0.00 | 0.0 |
| Metalaxyl (L) | 0.080 | 0.01 | 0.2 |
| Methamidophos | 0.001 | 0.07 | 73.9 |
| Methomyl | 0.003 | 0.00 | 0.4 |
| Pyraclostrobin | 0.030 | 0.02 | 0.7 |
| Thiamethoxam | 0.026 | 0.02 | 0.8 |

methomyl > boscalid > carbendazim imidacloprid > propiconazole. The results show a potential health risk associated mainly with methamidophos and cypermethrin in leafy vegetables in Chile. The HQs were summed up and the cHI of all residues was 135%, which was above 100%, including methamidophos and therefore should be considered a risk for consumers. When methamidophos was excluded from the risk assessment, cHI was 61%, which was below 100% and therefore not considered a health risk. Chen et al. (2011) found no health risk for consumers in fruits and vegetables, although the highest detected cypermethrin level (13.92 mg kg⁻¹ in cabbage) resulted in an intake that was determined to be 400% of the Acute Reference Dose (ARfD) value of 40 µg kg⁻¹ bw day⁻¹ when the WHO/GEMS high consumption diet was used. Lozowicka et al. (2015) found cHI values of 6.89%, 12.57% and 12.73% for organochlorines (endosulfan and dicofol), organophosphorus compounds (triazophos, chlorpyrifos and dimethoate) and fungicidal azoles (triadimefon, tebuconazole, triamenol, flusilazole and prochloraz), respectively, in tomatoes and cucumbers from Kazakhstan.

Conclusions

This study showed the presence of pesticide residues alone or in combination in leafy vegetables in three important regions of Chile. Furthermore, 65% of the samples showed two or more active ingredients and 27% of them were above the MRLs. Fungicides and insecticides were the most common pesticide types detected. The results provide important information on the current contamination status

of leafy vegetables in Chile. The results showed that the population was not exposed to health risks with the consumption of leafy vegetables, except for methamidophos, of which high levels could be considered a potential risk. Therefore, it is advisable to modify pest control management to reduce the use of methamidophos containing products because of their potential effect on human health.

Acknowledgement

This work was financed by the Ministry of Agriculture of Chile (Project 502061-70).

Disclosure statement

No potential conflict of interest was reported by the authors.

Funding

This work was financed by the Ministry of Agriculture of Chile [Project 502061-70 and Project 502187-70].

References

- Akoto O, Andoh H, Darko G, Eshun K, Osei-Fosu P. 2013. Health risk assessment of pesticides residue in maize and cowpea from Ejura, Ghana. *Chemosphere*. 92:67–73.
- Akoto O, Gavor S, Appah MK, Apau J. 2015. Estimation of human health risk associated with the consumption of pesticide-contaminated vegetables from Kumasi, Ghana. *Environ Monit Assess*. 187:244.
- Anastassiades M, Lehota SJ, Stajnbaher D, Schenck FJ. 2003. Fast and easy multiresidue method employing acetonitrile extraction/partitioning and “dispersive solid-phase extraction” for the determination of pesticide residue in produce. *J AOAC Int*. 86:412–431.
- Bempah CK, Buah-Kwofie A, Enimil E, Blewu B, Agyei-Martey G. 2012. Residues of organochlorine pesticides in vegetables marketed in Greater Accra Region of Ghana. *Food Control*. 25:537–542.
- Bhanti M, Taneja A. 2005. Monitoring of organochlorine pesticide residues in summer and winter vegetables from Agra, India - a case study. *Environ Monit Assess*. 110:341–346.
- Chen C, Qian Y, Chen Q, Tao C, Li C, Li Y. 2011. Evaluation of pesticide residues in fruits and vegetables from Xiamen, China. *Food Control*. 22:1114–1120.
- Chen C, Qian Y, Liu X, Tao C, Liang Y, Li Y. 2012. Risk assessment of chlorpyrifos on rice and cabbage in China. *Regul Toxicol Pharm*. 62:125–130.
- CODEx Alimentarius Commission 2011. [Internet]. [cited 2016 April 15]. Available from: <http://www.fao.org/fao-who-codexalimentarius/standards/list-of-standards/en/Accessed 15.4.16>.
- Darko G, Akoto O. 2008. Dietary intake of organophosphorus pesticide residues through vegetables from Kumasi, Ghana. *Food Chem Toxicol*. 46:3703–3706.
- De Putter H, Van Sauers-Muller A. (2008). Eindverslag Surituin deelproject: Terugdringen van residuu nrmen overschrijdingen in Surinaamse export groeten, 39 p. Available from: <http://www.wur.nl/en/Publication-details.htm?publicationId=publication-way-333831383637>
- Engler A, Nahuelhual L, Cofré G, Barrena J. 2012. How far from harmonization are sanitary, phytosanitary and quality-related standards? An exporter's perception approach. *Food Policy*. 37:162–170.
- ENS Chile. 2009-2010. Encuesta nacional de Salud. Ministerio de Salud, Gobierno de Chile. [Internet]. [cited 2016 April 10]. Available from: <http://web.minsal.cl/portal/url/item/bcb03d7bc28b64dfe040010165012d23.pdf>
- Esturk O, Yakar Y, Ayhan Z. 2011. Pesticide residue analysis in parsley, lettuce and spinach by LC-MS/MS. *J Food Sci Technol*. 51:458–466.
- FAO/WHO, 1997. Guidelines for predicting dietary intake of pesticide residue; Codex Alimentarius Commission. FAO/WHO Food Standards. [Internet]. [cited 2016 April 5]. Available from: http://www.who.int/foodsafety/publications/chem/en/pesticide_en.pdf
- Freeman S, Kaufman-Shriqui V, Berman T, Varsano R, Shahar DRManor O. 2016. Children's diets, pesticide uptake, and implications for risk assessment: an israeli case study. *Food Chem Toxicol*. 87:88–96.
- Gad Alla SA, Loutfy NM, Shendy AH, Ahmed MT. 2015. Hazard index, a tool for a long term risk assessment of pesticide residues in some commodities, a pilot study. *Regul Toxicol Pharm*. 73:985–991.
- Glat R. 2010. Scaling up political ecology: the case of non-authorized pesticides on fresh vegetables imported into the United States, 1996-2006. *Ann Assoc Am Geogr*. 100:327–355.
- Goh KT, Yew FS, Ong KH, Tan IK. 1990. Acute organophosphorus food poisoning caused by contaminated green leafy vegetables. *Arch Environ Health*. 45:180–184.
- González-Rodríguez RM, Rial-Otero R, Cancho-Grande B, Simal-Gándara J. 2008. Occurrence of fungicide and insecticide residues in trade samples of leafy vegetables. *Food Chem*. 107:1342–1347.
- Handsusch C, Wollni M, Villalobos P. 2013. Adoption of food safety and quality standards among Chilean raspberry producers – do smallholders benefits. *Food Policy*. 40:64–73.
- Hernández AF, Parrón T, Tsatsakis AM, Requena M, Alarcón R, López-Guarnido O. 2013. Toxic effects of pesticide mixtures at a molecular level: their relevance to human health. *Toxicology*. 307:136–145.
- INE, 2002, 2016. National Institute of Statistics in Chile. [Internet]. [cited 2016 April 1]. Available from: <http://www.ine.cl>
- Ivey KL, Hodgson JM, Croft KD, Lewis JR, Prince RL. 2015. Flavonoid intake and all-cause mortality. *Am J Clin Nutr*. 1:1012–1020.
- Jardim ANO, Mello DC, Goes FCS, Frota EF, Caldas ED. 2014. Pesticide residues in cashew apple, guava, kaki and peach: GC- μ ECD, GC-PPD and LC-MS/MS multiresidue method validation, analysis and cumulative acute risk assessment. *Food Chem*. 164:195–204.
- Köck-Schulmeyer M, Villagrasa M, López De Alda M, Céspedes-Sánchez R, Ventura F, Barceló D. 2013. Occurrence and behavior of pesticides in wastewater treatment plants and their environmental impact. *Sci Total Environ*. 458:466–476.

- Lemos J, Sampedro MC, Ariño A, Ortiz A, Barrio RJ. 2016. Risk assessment of exposure to pesticides through dietary intake of vegetables typical of the Mediterranean diet in the Basque Country. *J Food Compost Anal.* 49:35–41.
- Li R, He L, Wei W, Hao L, Ji X, Zhou Y, Wang Q. 2015. Chlorpyrifos residue levels on field crops (rice, maize and soybean) in China and their dietary risks to consumers. *Food Control.* 51:212–217.
- Li W, Tai L, Liu J, Gai Z, Ding G. 2014. Monitoring of pesticide residues levels in fresh vegetable from Hebei Province, North China. *Environ Monit Assess.* 186:6341–6349.
- Lozowicka B, Abzeitova E, Sagitov A, Kaczynski P, Toleubayev K, Li A. 2015. Studies of pesticide residues in tomatoes and cucumbers from Kazakhstan and the associated health risks. *Environ Monit Assess.* 187:609.
- McEwen FL, Ritcey GM, Braun H, Frank R, Ripley BD. 1980. Foliar pesticide residues in relation to worker re-entry. *Pestic Sci.* 11:643–650.
- Melo O, Engler A, Nahuehual L, Cofre G, Barrera J, Sanitary D. 2014. Phytosanitary, and quality-related standards affect international trade? Evidence from Chilean fruit exports. *World Dev.* 54:350–359.
- Oerke EC, Dehne HW. 2004. Safeguarding production - losses in major crops and the role of crop protection. *Crop Prot.* 23:275–285.
- Park DW, Kim KG, Choi EA, Kang GR, Sun T, Yang Y, Moon S, Ryong H, Kim S, Cho B. 2016. Pesticide residues in leafy vegetables, stalk and stem vegetables from South Korea: a long-term study on safety and health risk assessment. *Food Addit Contam.* 33:105–118.
- Parrón T, Requena M, Hernández AF, Alarcón R. 2014. Environmental exposure to pesticides and cancer risk in multiple human organ systems. *Toxicol Lett.* 230:157–165.
- Payá P, Anastassiades M, Mack D, Sigalova I, Tsdelen B, Oliva J, Barba A. 2007. Analysis of pesticide residues using the Quick Easy Cheap Effective Rugged and Safe (QuEChERS) pesticide multiresidue method in combination with gas and liquid chromatography and tandem mass spectrometric detection. *Anal Bioanal Chem.* 389:1697–1714.
- Perz RC, Van Lishaut H, Schwack W. 2000. CS₂ blinds in Brassica crops: false positive results in the dithiocarbamate residue analysis by the acid digestion method. *J Ag Food Chem.* 48:792–796.
- SANTE, European Commission, Directorate General for Health and Food Safety, 2015. Guidance document on analytical quality control and method validation procedures for pesticides residues analysis in food and feed. SANTE/11945/2015, https://ec.europa.eu/food/sites/food/files/plant/docs/pesticides_mrl_guidelines_wrkdoc_11945.pdf [assessed April 2016].
- Sapbamrer R, Hongsibsong S. 2014. Organophosphorus pesticide residues in vegetables from farms, markets, and a supermarket around Kwan Phayao Lake of Northern Thailand. *Arch Environ Contam Toxicol.* 67:60–67.
- Srivastava AK, Trivedi P, Srivastava MK, Lohani MSrivastava LP. 2011. Monitoring of pesticide residues in market basket samples of vegetable from lucknow city, india: quechers method. *Environ Monit Assess.* 176:465–472.
- Szpyrka E, Kurdziel A, Matyaszek A, Podbielska M, Rupa J, Słowik-Borowiec M. 2015. Evaluation of pesticide residues in fruits and vegetables from the region of south-eastern Poland. *Food Control.* 48:137–142.
- Wang X, Zhang H, Xu H, Qi P, Ji X, Wang Q, Wang X. 2013. Direct chiral determination of acephate and its metabolite methamidophos in vegetables using QuEChERS by gas chromatography-tandem mass spectrometry. *Food Anal Methods.* 6:133–140.
- WHO/FAO. (2013). *Codex Alimentarius Commission, Procedural manual.* 2013 (20th ed) Rome. [Internet]. [cited 2016 Feb 10]. <http://www.fao.org/3/a-i3243e.pdf>
- Yu R, Liu Q, Liu J, Wang Q, Wang Y. 2016. Concentrations of organophosphorus pesticides in fresh vegetables and related human health risk assessment in Changchun, Northeast China. *Food Control.* 60:353–360.